Brain Research, 573 (1992) 356–357
Elsevier

Author Index

Adachi, S., Oka, J.-I., Nagao, T. and Fukuda, H., Cyclic AMP analog activates Na^+-dependent inward currents in dissociated frog motoneurons, 349
Andreasen, M., see Little, H.J., 243
Angst, M.J., see Sandner, G., 276
Anwyl, R., see Oonnor, J.J., 190
Artola, A., see Bröcher, S., 27

Bach, F., Flügge, G. and Wuttke, W., GABAergic influence on the development of the sexually dimorphic nucleus of male and female rats, 341
Baudry, M., see Tocco, G., 228
Baum, L., Masliah, E., Iimoto, D.S., Hansen, L.A., Halliday, W.C. and Saitoh, T., Casein kinase II is associated with neurofibrillary tangles but is not an intrinsic component of paired helical filaments, 126
Blessing, W.W., see Li, Y.-W., 147
Bluthé, R.-M., Dantzer, R. and Kelley, K.W., Effects of interleukin-1 receptor antagonist on the behavioral effects of lipopolysaccharide in rat, 318
Bodenant, C., see Laquerrière, A., 251
Briand, P.-A., see Eder-Colli, L., 284
Bridges, R.S., see Felicio, L.F., 133
Briggs, C.A., Potentiation of nicotinic transmission in the rat superior cervical sympathetic ganglion: effects of cyclic GMP and nitric oxide generators, 139
Bröcher, S., Artola, A. and Singer, W., Agonists of cholinergic and noradrenergic receptors facilitate synergistically the induction of long-term potentiation in slices of rat visual cortex, 27

Calvet, M.-C., see Levallois, C., 327
Chapman, V. and Dickenson, A.H., The combination of NMDA antagonism and morphine produces profound antinociception in the rat dorsal horn, 321
Chayvialle, J.-A., see Mick, G., 311
Chiappinelli, V.A., see Nooney, J.M., 77
Ciofi, P., see Maubert, E., 153
Clincke, G., see De Ryck, M., 44

Cox, B.M., see Izenwasser, S., 119
Cordisco, B.R., see Griffith, J.K., 1
Cronin, J., Obenaus, A., Houser, C.R. and Dudek, F.E., Electrophysiology of dentate granule cells after kainate-induced synaptic reorganization of the mossy fibers, 305
Crowley, W.R., see Sahu, A., 235

Dantzer, R., see Bluthé, R.-M., 318
De Ryck, M., Van Reempts, J., Duytschaever, H., Van Deuren, B. and Clincke, G., Neocortical localization of tactile/proprioceptive limb placing reactions in the rat, 44
Deacon, T.W., Cortical connections of the inferior arcuate sulcus cortex in the macaque brain , 8
Di Scala, G., see Sandner, G., 276
Dickenson, A.H., see Chapman, V., 321
Ding, Z.-Q., see Li, Y.-W., 147
Dora, E., Hines, K., Kunos, G. and McLaughlin, A.C., Significance of an opiate mechanism in the adjustment of cerebrocortical oxygen consumption and blood flow during hypercapnic stress, 293
Dudek, F.E., see Cronin, J., 305
Dunant, Y., see Eder-Colli, L., 284
Dupouy, J.-P., see Maubert, E., 153
Duytschaever, H., see De Ryck, M., 44

Eder-Colli, L., Briand, P.-A. and Dunant, Y., Membrane-bound choline acetyltransferase of the torpedo has characteristics of an integral membrane protein and can be solubilized by proteolysis, 284
Eysel, U.T., see Funke, K., 217

Felicio, L.F. and Bridges, R.S., Domperidone induces a probenecid-sensitive rise in immunoreactive prolactin in cerebroventricular perfusates in female rats, 133
Flügge, G., see Bach, F., 341
Fowler, J.C., Escape from inhibition of synaptic transmission during in vitro hypoxia and hypoglycemia in the hippocampus, 169
Fukuda, H., see Adachi, S., 349
Funke, K. and Eysel, U.T., EEG-dependent modulation of response dynamics of cat dLGN relay cells and the contribution of corticogeniculate feedback, 217
Fuxe, K., see Oonnor, W.T., 209

Girard, M., see Mick, G., 311
Goldman, R.D., see Yang, H.-Y., 161
Gonzalez, B., see Laquerrière, A., 251
Griffith, J.K., Cordisco, B.R., Lin, C.-W. and LaManna, J.C., Distribution of intracellular pH in the rat brain cortex after global ischemia as measured by color film histophotometry of Neutral red, 1
Grill, H.J. and Schwartz, G.J., The contribution of gustatory nerve input to oral motor behavior and intake-based preference. II. Effects of combined chorda tympani and glossopharyngeal nerve section in the rat, 105
Grill, H.J., Schwartz, G.J. and Travers, J.B., The contribution of gustatory nerve input to oral motor behavior and intake-based preference. I. Effects of chorda tympani or glossopharyngeal nerve section in the rat, 95
Gruol, D.L., see Yool, A.J., 83

Halliday, W.C., see Baum, L., 126
Hansen, L.A., see Baum, L., 126
Hatase, O., see Minamoto, Y., 345
Heller, H.C., see Prosser, R.A., 336
Hines, K., see Dora, E., 293
Hosokawa, K., see Minamoto, Y., 345
Houser, C.R., see Cronin, J., 305
Hyde, J.F., Effects of phencyclidine on 5-hydroxytryptophan- and suckling-induced prolactin release, 204

Iimoto, D.S., see Baum, L., 126
Itano, T., see Minamoto, Y., 345
Izenwasser, S. and Cox, B.M., Inhibition of dopamine uptake by cocaine and nicotine: tolerance to chronic treatments, 119

Janjua, N.A., see Minamoto, Y., 345
Johansson, B.B., see Westergren, I., 324
Johnson-Seaton, D., see Yang, H.-Y., 161

Kabai, P., Kovach, J. and Vadasz, C., Neural correlates of genetically determined and acquired color preferences in quail chicks, 260
Kalra, P.S., see Sahu, A., 235
Kalra, S.P., see Sahu, A., 235
Kamenka, J.-M., see Levallois, C., 327
Kelley, K.W., see Bluthé, R.-M., 318
Komiyama, A. and Suzuki, K., Age-related differences in proliferative responses of Schwann cells during Wallerian degeneration, 267
Kovach, J., see Kabai, P., 260
Krantic, S., Robitaille, Y. and Quirion, R., Deficits in the somatostatin SS_1 receptor sub-type in frontal and temporal cortices in Alzheimer's disease, 299
Krieger, R.M., see Yool, A.J., 83
Kunishita, T., see Satoh, J.-i., 37
Kunos, G., see Dora, E., 293

LaManna, J.C., see Griffith, J.K., 1
Lambert, J.D.C., see Little, H.J., 243
Lambert, J.J., see Nooney, J.M., 77
Langub Jr., M.C. and Watson Jr., R.E., Estrogen receptive neurons in the preoptic area of the rat are postsynaptic targets of a sexually dimorphic enkephalinergic fiber plexus, 61
Laquerrière, A., Leroux, P., Gonzalez, B., Bodenant, C., Tayot, J. and Vaudry, H., Somatostatin receptors in the human cerebellum during development, 251
Larsen, P.J., Sheikh, S.P. and Mikkelsen, J.D., Osmotic regulation of neuropeptide Y and its binding sites in the magnocellular hypothalamo-neurohypophysial pathway, 181
Larson, G., see Mayfield, R.D., 331
Leroux, P., see Laquerrière, A., 251
Levallois, C., Calvet, M.-C., Kamenka, J.-M., Petite, D. and Privat, A., TCP enhances the survival of human fetal

spinal cord cells in culture, 327
Li, Y.-W., Ding, Z.-Q., Wesselingh, S.L. and Blessing, W.W., Renal and adrenal sympathetic preganglionic neurons in rabbit spinal cord: tracing with Herpes simplex virus, 147
Lieska, N., see Yang, H.-Y., 161
Lin, C.-W., see Griffith, J.K., 1
Little, H.J., Andreasen, M. and Lambert, J.D.C., Chronic treatment with a benzodiazepine agonist in vivo increases the actions of the benzodiazepine partial inverse agonist, FG7142, on the hippocampal slice in vitro, 243

Maelicke, A., see Reuss, S., 114
Maren, S., see Tocco, G., 228
Masliah, E., see Baum, L., 126
Matsui, H., see Minamoto, Y., 345
Maubert, E., Ciofi, P., Tramu, G., Mazzuca, M. and Dupouy, J.-P., Early transient expression of somatostatin (SRIF) immunoreactivity in dorsal root ganglia during ontogenesis in the rat , 153
Mayfield, R.D., Larson, G. and Zahniser, N.R., Cocaine-induced behavioral sensitization and D_1 dopamine receptor function in rat nucleus accumbens and striatum, 331
Mazzuca, M., see Maubert, E., 153
McLaughlin, A.C., see Dora, E., 293
Mick, G., Najimi, M., Girard, M. and Chayvialle, J.-A., Evidence for a substance P containing subpopulation in the primate suprachiasmatic nucleus, 311
Mikkelsen, J.D., see Larsen, P.J., 181
Miller, J.D., see Prosser, R.A., 336
Minamoto, Y., Itano, T., Tokuda, M., Matsui, H., Janjua, N.A., Hosokawa, K., Okada, Y., Murakami, T.H., Negi, T. and Hatase, O., In vivo microdialysis of amino acid neurotransmitters in the hippocampus in amygdaloid kindled rat, 345
Murakami, T.H., see Minamoto, Y., 345

Nagao, T., see Adachi, S., 349
Najimi, M., see Mick, G., 311
Negi, T., see Minamoto, Y., 345
Nemeroff, C.B., see Owens, M.J., 353
Nooney, J.M., Lambert, J.J. and Chiappinelli, V.A., The interaction of κ-bungarotoxin with the nicotinic receptor of bovine chromaffin cells, 77

Obenaus, A., see Cronin, J., 305
Oonnor, J.J., Rowan, M.J. and Anwyl, R., Serotoninergic depression of auditory evoked responses recorded in the rat hippocampus: effect of repeated buspirone treatment, 190
Oonnor, W.T., Tanganelli, S., Ungerstedt, U. and Fuxe, K., The effects of neurotensin on GABA and acetylcholine release in the dorsal striatum of the rat: an in vivo mirodialysis study, 209
Oka, J.-I., see Adachi, S., 349
Okada, Y., see Sakurai, T., 197
Okada, Y., see Minamoto, Y., 345
Owens, M.J., Ritchie, J.C. and Nemeroff, C.B., 5α-Pregnane-3α,21-diol-20-one (THDOC) attenuates mild stress-induced increases in plasma corticosterone via a non-glucocorticoid mechanism: comparison with alprazolam, 353

Petite, D., see Levallois, C., 327
Privat, A., see Levallois, C., 327
Prosser, R.A., Heller, H.C. and Miller, J.D., Serotonergic phase shifts of the mammalian circadian clock: effects of tetrodotoxin and high Mg, 336

Quirion, R., see Krantic, S., 299

Reuss, S., Schröder, B., Schröder, H. and Maelicke, A., Nicotinic cholinoceptors in the rat pineal gland as analyzed by Western blot, light- and electron microscopy, 114
Ritchie, J.C., see Owens, M.J., 353
Robitaille, Y., see Krantic, S., 299
Rocha, B., see Sandner, G., 276
Rowan, M.J., see Oonnor, J.J., 190

Sahu, A., Crowley, W.R., Kalra, P.S. and Kalra, S.P., A selective sexually dimorphic response in the median eminence neuropeptide Y* , 235
Saitoh, T., see Baum, L., 126
Sakurai, T. and Okada, Y., Selective reduction of glutamate in the rat superior colliculus and dorsal lateral geniculate nucleus after contralateral enucleation, 197
Sandner, G., Di Scala, G., Rocha, B. and Angst, M.J., *C-fos* immunoreactivity in the brain following unilateral electrical stimulation of the dorsal periaqueductal gray in freely moving rats, 276
Satoh, J.-i., Yamamura, T., Kunishita, T. and Tabira, T., Heterogeneous induction of 72-kDa heat shock protein (HSP72) in cultured mouse oligodendrocytes and astrocytes, 37
Schröder, B., see Reuss, S., 114
Schröder, H., see Reuss, S., 114
Schwartz, G.J., see Grill, H.J., 105
Schwartz, G.J., see Grill, H.J., 95
Sheikh, S.P., see Larsen, P.J., 181
Shors, T.J., see Tocco, G., 228
Singer, W., see Bröcher, S., 27
Somjen, G.G., see Young, J.N., 70
Surchev, L., Freeze-etched postsynaptic membranes in the visual cortex reveal different types of synapses including mixed synapses, 174
Suzuki, K., see Komiyama, A., 267

Tabira, T., see Satoh, J.-i., 37
Tanganelli, S., see Oonnor, W.T., 209
Tayot, J., see Laquerrière, A., 251
Thompson, R.F., see Tocco, G., 228
Tocco, G., Maren, S., Shors, T.J., Baudry, M. and Thompson, R.F., Long-term potentiation is associated with increased [^{3}H]AMPA binding in rat hippocampus, 228
Tokuda, M., see Minamoto, Y., 345
Tramu, G., see Maubert, E., 153
Travers, J.B., see Grill, H.J., 95

Ungerstedt, U., see Oonnor, W.T., 209

Vadasz, C., see Kabai, P., 260
Van Deuren, B., see De Ryck, M., 44
Van Reempts, J., see De Ryck, M., 44
Vaudry, H., see Laquerrière, A., 251

Watson Jr., R.E., see Langub Jr., M.C., 61
Wesselingh, S.L., see Li, Y.-W., 147
Westergren, I. and Johansson, B.B., NBQX, an AMPA antagonist, reduces glutamate-mediated brain edema, 324
Wuttke, W., see Bach, F., 341

Yamamura, T., see Satoh, J.-i., 37
Yang, H.-Y., Lieska, N., Goldman, R.D. and Johnson-Seaton, D., Distinct developmental subtypes of cultured non-stellate rat astrocytes distinguished by a new glial intermediate filament-associated protein, 161
Yool, A.J., Krieger, R.M. and Gruol, D.L., Multiple ionic mechanisms are activated by the potent agonist quisqualate in cultured cerebellar Purkinje neurons , 83
Young, J.N. and Somjen, G.G., Suppression of presynaptic calcium currents by hypoxia in hippocampal tissue slices, 70

Zahniser, N.R., see Mayfield, R.D., 331